

There is no evidence that reduction mammoplasty causes breast cancer. Nor does it prevent your breast from being examined for cancer in the usual way. Breast tissue removed at reduction is sent to histology for testing.

**What should you do before your operation?**

Your surgeon is likely to recommend that you reduce weight if you are overweight and to make alternative arrangements if you are on the contraceptive pill.

**What can you expect at the time of the operation?**

This procedure is carried out under a general anaesthetic and when you wake at the end of the operation there will be some discomfort which will last for two or three days. You will be given suitable pain killing injections or tablets. You may need to remain in hospital for two days. Most stitches are dissolvable but occasionally may be removed between ten to fourteen days. You will be tired and require help at home for a period of time, from two to six weeks depending on your age and general fitness. A well fitting bra will need to be worn following surgery but because of the postoperative swelling the final size of your breasts may not be obvious for several weeks. There is likely to be some tenderness and lumpiness of the breasts for several weeks or even months following surgery but there is no reason why you cannot sunbathe and go swimming once the scars have fully healed.

© J D Frame

September 2019

Consultations at:  
Springfield Hospital,  
Chelmsford, Essex

Secretary:-  
Springfield Hospital  
Lawn Lane  
Springfield  
Chelmsford  
Essex  
CM1 7GU

Phone: 01245 460981  
Fax: 01245 460991  
Email: [info@professorjamesframe.co.uk](mailto:info@professorjamesframe.co.uk)  
[www.professorjamesframe.co.uk](http://www.professorjamesframe.co.uk)  
[www.ukaaps.org](http://www.ukaaps.org)

**Professor J D Frame**  
FRCS, FRCS (Plast.)

# Breast Reduction

**Professor J D Frame**  
FRCS, FRCS (Plast.)

Consultant Plastic  
and  
Aesthetic Surgeon

## Breast Reduction

(Reduction Mammoplasty)

Tel: 01245 460981

# Breast Reduction (Reduction Mammoplasty)

The size of women's breasts may be determined by several factors, such as inherited genes, body weight and hormonal influences. They can, therefore, be a problem to some women early in adolescence or may not become uncomfortable until middle age following the menopause or the use of HRT. The problem of large breasts, however, may cause similar problems at all ages and these are chiefly backache, neck pain, grooves in the shoulders from bra straps, rashes under the breasts and the feeling of self-consciousness and embarrassment. Because of the sexual nature of breasts the undue prominence may attract unwanted attention from the opposite sex, comments and sexual innuendoes. These can cause psychological distress to many women. One of the commonest complaints of women with large breasts is that it is very difficult to wear fashionable clothes and indulge in active sports, particularly in the summer months.

## What can be done?

Reduction mammoplasty is an operation which removes the excess fat and skin from the breasts, which are re-shaped and the nipples repositioned to form smaller breasts. All of them will involve a scar around the areola of the breast but there will be vertical and perhaps horizontal scars as well. Different scars will have advantages and disadvantages and your surgeon may select, with your approval, the best technique in your particular case. Reduction mammoplasty may be used to correct asymmetry of the breast, where one breast is very much larger than the other. Sometimes, marginal volume difference can be improved by liposuction alone.

## What are the consequences?


Apart from the change of shape and reduction in size the most obvious consequences are the scars. These are designed to be hidden whilst wearing normal clothing and as far as possible are designed to lie under the average bra or bikini top. Over the months following surgery the scars will fade from being red, possibly thick and uncomfortable, to becoming much more pale and less obvious. However, they will always be present and visible when clothing is not worn and the scars will vary from one woman to another. In some they may be very thin, in others they may stretch and become quite red and possibly ugly. In the vast majority of women, however, the scars are acceptable and a small trade off for the benefit of dealing with the problems of large breasts. Very few women are able to breast feed following breast reduction surgery, as some of the milk ducts are separated from the underlying glands and at the time of pregnancy the milk supply will gradually dry up, sometimes with the assistance of hormone treatment. Breast reduction is no contra indication to pregnancy but young women may well wish to take the

fact that they are unable to breast feed into account before embarking on this procedure.


The nipples are likely to be very much less sensitive following surgery due to the nature of the cuts and the nerve supply and it is quite possible that numbness will extend over part of the breast as well. Sometimes the nipples become hyper-sensitive, if and when they recover sensation.

## Will the improvement last?

Unless your operation is done at an age when your breasts are still



Before operation



Vertical Scar Reduction

Inverted T Scar

growing, they should not regrow afterwards. They will, however, increase in size if you either put on weight or become pregnant and decrease in size if you lose weight. Even normal breasts have a tendency to droop with time and you can expect some change in shape to occur after a reduction mammoplasty. You can delay this tendency by supporting your breasts in well fitting bras. The heavier the remaining breast tissue, the more likely to drop at a later stage.

## What are the limitations?

It is very important that you discuss thoroughly with your surgeon beforehand the size of breasts you wish to achieve. Many women wish for a very radical reduction but in order to achieve this the shape and aesthetic quality of the breasts may be compromised. It

is also possible that there may be a degree of asymmetry. This is often less than existed before the surgery. In women with very large breasts the benefits of a significant reduction may outweigh the potential imperfections of poor shape and loss of nipple function and sensitivity. However, in women with breasts which are only slightly larger than normal very careful thought must be given to the scarring and the potential shape and size as the outcome of the surgery may not be as aesthetically pleasing as a normal breast appearance.

## What are the risks?

Any major operation with a general anaesthetic carries a small risk of chest infection particularly among people who smoke and there is also a small risk of thrombosis in the veins of the leg, particularly for patients who are taking the contraceptive pill. Long haul flights in the post-op period should be delayed for at least six weeks.

Occasionally, bleeding can occur after the operation is finished. This may require surgical intervention. Infection from germs harbouring in the ducts of the breast can be troublesome. This can be treated with antibiotics but it will delay the healing process. Scars are likely to be worse to start with and there may be a need to re-stitch them at a later date. If you have a discharge from your nipple, it is most important to tell your surgeon about it before your operation.

Occasionally, skin can become sloughy and form a scab which gradually separates to leave a broad scar. The nipple disc and the skin where the scar meets underneath the breast are parts most likely to be affected. People who smoke are at greater risk of this happening. Usually the scars settle well to end up as white lines but they will always be noticeable. However, some people have an inborn tendency for scars to stretch and sometimes they can stay thick, red and irritable for a long time.

When reducing large breasts using the vertical scar technique, it may occasionally be necessary to adjust the folds of skin at the end of the scar, both between the breasts and at the sides. This can simply be carried out under local anaesthetic several months later. It is unlikely for this to be needed if the Inverted T scar is used.

Secretary:-  
Springfield Hospital  
Lawn Lane  
Springfield  
Chelmsford  
Essex

Phone: 01245 460981

Fax: 01245 460991

Email: [info@professorjamesframe.co.uk](mailto:info@professorjamesframe.co.uk)